

The Polyakov loop and the heat kernel expansion at finite temperature

E. Megías, E. Ruiz Arriola, and L.L. Salcedo

Departamento de Física Moderna, Universidad de Granada, E-18071 Granada, Spain

(Dated: February 1, 2008)

The lower order terms of the heat kernel expansion at coincident points are computed in the context of finite temperature quantum field theory for flat space-time and in the presence of general gauge and scalar fields which may be non Abelian and non stationary. The computation is carried out in the imaginary time formalism and the result is fully consistent with invariance under topologically large and small gauge transformations. The Polyakov loop is shown to play a fundamental role.

PACS numbers: 11.10.Wx, 11.10.Kk, 11.15.-q, 11.10.Jj

Keywords: Finite temperature; heat kernel expansion; Effective action; Gauge invariance

The heat kernel $e^{-\tau K}$, where K is an Euclidean Klein-Gordon operator, was introduced by Schwinger [1] in quantum field theory as a tool to regularize ultraviolet divergences in a gauge invariant way. The procedure of systematically isolating these divergences was developed by DeWitt through an asymptotic expansion in powers of the proper time τ [2, 3]. The heat kernel as well as its expansion has also been applied to study spectral densities and indices of Dirac operators ($\hat{D}$) [4, 5] in terms of effective Klein-Gordon operators ($\hat{D}^\dagger \hat{D}$), to compute the ζ -function [6] and anomalies of such operators [7], to properly define the effective action of chiral gauge theories [8], to the Casimir effect [9], etc. Exact calculations of the heat kernel at coincident points are available in particular manifolds [10, 11], and the coefficients of the expansion have been computed to rather high orders in several setups, including curved spaces with and without boundary, and in presence of non Abelian gauge fields and non Abelian scalar fields, using different methods [8, 12, 13, 14, 15, 16, 17]. In this work we deal with the extension of these results to finite temperature in flat space-time.

As is well known, in the Matsubara formalism for quantum field theory at finite temperature, wave functions are required to be periodic for bosons and antiperiodic for fermions, with period β (the inverse temperature) and the external fields are also periodic. This introduces a compactified Euclidean time [18]. The heat kernel can be regarded as the evolution operator (in imaginary proper time τ) of a first quantized particle [1], thus it admits a path integral representation, which at finite temperature must include also multiply connected paths in the compactified space-time. The standard results quoted e.g. in [8] for the heat kernel expansion refer to general smooth manifolds without boundary and thus, in principle, they would directly apply to finite temperature as well. However, the standard formulas are not sensitive to the compactification of time; they only sum paths which are topologically trivial (contractible to a point). Actually this correctly reproduces the analytical part in a strict expansion in powers of the proper time, but neglects the exponentially suppressed finite temperature contributions

[19]. To see any temperature effect in the heat kernel expansion it is necessary to include also the paths that wind once or more around the compactified time. This is automatically achieved in the canonical formalism by replacing the continuous energy variable by the discrete Matsubara frequencies. More generally, as already argued in [20] (see also [21]), in the presence of gauge fields, finite temperature effects are tied to the presence of the (untraced) Polyakov loop

$$\Omega(x) = T \exp \left(- \int_{x_0}^{x_0+\beta} A_0(x'_0, \mathbf{x}) dx'_0 \right) \quad (1)$$

(T refers to temporal ordering, this is required to allow for the non Abelian case). The presence of $\Omega(x)$ is to be expected on general grounds. Let us regard the heat kernel at finite temperature as the evolution operator of a first quantized particle. The space-time manifold defines a d -dimensional cylinder which plays of role of space for the particle and $-i\tau$ is the time. If the circulation of the “vector potential” A_μ around the cylinder is non vanishing this signals the presence of an Aharonov-Bohm magnetic flux through its interior, which will affect the propagation. The particle will pick up a phase $\Omega(x)$ after looping once around the cylinder (in the general case a non Abelian phase [22]).

A computation of the thermal heat kernel coefficients was presented in [23] where a common renormalization for all the coefficients, $a_n^T = f(\tau/\beta^2) a_n$, was found. However, because the temperature effects account for the fact that Euclidean time is compact, an explicit breaking of Lorentz invariance is expected and the previous factorization cannot hold in general (see e.g. [24]). The former computation refers to flat space-time (without boundaries) and non Abelian scalar and gauge fields. In [25] a similar study is presented where curvature is allowed but fields are assumed to be static. These authors find that $f(\tau/\beta^2) a_n$ is augmented with new terms that break Lorentz symmetry, so-called anomalous terms also found in [26]. However both calculations are incomplete since they are in conflict with results for Abelian and stationary configurations (in flat space) presented in [24]. There

it is shown that the Euclidean scalar potential $A_0(\mathbf{x})$ can appear in the heat kernel (outside covariant derivatives) provided that it does so as a periodic variable with period $2\pi/\beta$, to be consistent with (large) gauge invariance. This is a particular manifestation of the relevance of the Polyakov loop.

We present here the first systematic calculation of the heat kernel expansion at finite temperature at coincident points (in flat space without boundaries) for general backgrounds which may be non Abelian and non stationary. The result is gauge covariant and depends locally on the covariant derivatives and the Polyakov loop. The Klein-Gordon operator to be considered is of the type $K = M - D_\mu^2$, where $D_\mu = \partial_\mu + A_\mu(x)$ is the gauge covariant derivative and $M(x)$ is a Lorentz scalar. Both $A_\mu(x)$ and $M(x)$ are allowed to be matrices in some internal space (the treatment applies to Abelian and non Abelian theories). One can form local and gauge covariant operators by combining the basic operators M , $F_{\mu\nu} = [D_\mu, D_\nu]$ and their covariant derivatives, e.g. $\mathcal{O} = F_{\mu\nu} \mathcal{D}_\mu \mathcal{D}_\nu M$, where $\mathcal{D}_\mu X$ stands for $[D_\mu, X]$. The local operators can be classified according to their mass dimension. For each dimension there is a finite number of them. At zero temperature, under suitable regularity conditions, one can asymptotically expand an expression of the form $\langle x|f(K)|x \rangle$ as a sum of local operators, i.e.

$$\langle x|f(K)|x \rangle = \sum_r f_r \mathcal{O}_r(x), \quad (2)$$

$\mathcal{O}_r$ denotes a linear basis including all possible local operators and f_r are some numerical coefficients which depend on the function f . In particular, for the matrix elements of the heat kernel at coincident points one obtains the familiar Seeley-DeWitt asymptotic expansion

$$\langle x|e^{-\tau K}|x \rangle = (4\pi\tau)^{-d/2} \sum_{n=0}^{\infty} a_n(x) \tau^n, \quad (3)$$

d being the space-time dimension. The local operators $a_n(x)$ themselves do not depend on d or τ and have mass dimension $2n$. Of course, once one has the expansion for the heat kernel, the use of integral representations [1] allows to obtain the expansion for other functions $f(K)$, such as the ζ -function [3, 6], the effective action, the resolvent or the effective current, without a new computation.

At finite temperature, because the Euclidean time coordinate is compactified, one can expect to expand $\langle x|f(K)|x \rangle$ in terms of operators which are local regarding the spatial coordinates but not in the time coordinate [27]. Nevertheless, the following asymptotic expansion still holds [20]

$$\langle x|f(K)|x \rangle = \sum_r f_r(\Omega(x)) \mathcal{O}_r(x). \quad (4)$$

That is, at finite temperature the numerical coefficients f_r become functions of the Polyakov loop $\Omega(x)$. This result is remarkable on several accounts. First, although full space-time locality does not hold at finite temperature, still the result looks space-time local if written in terms of the field $\Omega(x)$ and the other local operators. (Note that $\Omega(x)$ is local regarding gauge transformations.) Second, full gauge covariance (the covariance under continuous local similarity transformations in internal space) is manifest, since this property is shared by the Polyakov loop and the local operators. This is not always automatically true for other expansions; because the space-time is not simply connected in the imaginary time formalism, in addition to the usual topologically small gauge transformations there are also large transformations (i.e., not reachable by composition of infinitesimal ones). Covariance under such large transformations fails to hold, for instance, in simple-minded perturbation theory [27, 28].

In the non Abelian case $\Omega(x)$ needs not commute with the other quantities. The fact that it appears to the left in (4) is a matter of choice (putting it to the right would yield a different set of functions f_r). We remark that the operators $\mathcal{O}_r$ are local and gauge covariant but not necessarily Lorentz invariant. That is, operators with different Lorentz indices may have different weights at finite temperature. Rotational invariance is of course preserved.

As we have argued above, for many purposes it is sufficient to consider the expansion for the heat kernel. At finite temperature the result can still be written as

$$\langle x|e^{-\tau K}|x \rangle = (4\pi\tau)^{-d/2} \sum_n a_n^T(x) \tau^n. \quad (5)$$

The thermal heat kernel coefficients a_n^T contain local operators, and depend on Ω , τ and β , but not on d .

At finite temperature the heat kernel expansion consists actually of three expansions. One in powers of M , another in powers of $\mathcal{D}_i$, and finally another expansion in powers of $\mathcal{D}_0$. The first two expansions are standard and no different to those done at zero temperature, but the expansion in $\mathcal{D}_0$ is tricky and has to be carried out with care. This is because a naive expansion in temporal derivatives is in conflict with gauge invariance. This is easy to understand by noting that the derivative expansion refers to external fields with very wide profiles. Whereas this is unproblematic in the spatial direction, the situation in the temporal direction is different since the Euclidean time is periodic and dilatations in the temporal direction break the periodicity condition.

For the calculation we use the method of symbols which is based on the identity [29, 30]

$$\langle x|f(D_\mu, M)|x \rangle = \frac{1}{\beta} \sum_{p_0} \int \frac{d^{d-1}p}{(2\pi)^{d-1}} \langle x|f(D_\mu + ip_\mu, M)|0 \rangle. \quad (6)$$

$f(D_\mu, M)$ is any (ultraviolet finite) operator constructed out of D_μ and M , and $|0\rangle$ is the zero momentum state, $\langle x|0\rangle = 1$. In practice this means that ∂_μ in D_μ vanishes when it reaches the right end, and the brackets $\langle x|\cdot|0\rangle$ are often not explicit. An application of the symbols method formula in the temporal direction only, gives

$$\begin{aligned}\langle x|e^{-\tau K}|x\rangle &= \frac{1}{\beta} \sum_{p_0} \langle x|e^{-\tau(M-Q^2-D_i^2)}|x\rangle, \\ Q &= ip_0 + D_0.\end{aligned}\quad (7)$$

For the spatial part we can directly use the known heat kernel expansion at zero temperature [8, 12] on the effective $(d-1)$ -dimensional Klein-Gordon operator $(M-Q^2)-D_i^2$. The operators Q are then moved to the left generating commutators of the form $[Q, \cdot]$. These commutators are just $D_0 = [D_0, \cdot]$ since p_0 is a c-number. The Q 's not in commutators, placed at the left of the expressions so obtained, come in the form $Q^n e^{\tau Q^2}$ (cf. (11)). The final step is to use the operator identity [20]

$$e^{-\beta D_0} = e^{-\beta \partial_0} \Omega(x). \quad (8)$$

Because all external fields are periodic, $e^{-\beta \partial_0} = 1$ and $e^{-\beta D_0}$ is equivalent to $\Omega(x)$. In this way the operator Q defined in (7) reduces to a function of x

$$Q = ip_0 - \frac{1}{\beta} \log(\Omega). \quad (9)$$

An explicit computation gives, up to operators of mass dimension four [37],

$$\begin{aligned}a_0^T &= \varphi_0 \\ a_1^T &= -\varphi_0 M \\ a_{3/2}^T &= \varphi_1 \left(M_0 - \frac{1}{3} E_{ii} \right) \\ a_2^T &= -\left(\frac{1}{2} \varphi_0 + \frac{2}{3} \varphi_2 \right) M_{00} \\ &\quad + \left(\frac{1}{6} \varphi_0 + \frac{1}{3} \varphi_2 \right) E_{0ii} + \left(\frac{1}{3} \varphi_0 + \frac{1}{3} \varphi_2 \right) E_i^2 \\ &\quad + \varphi_0 \left(\frac{1}{2} M^2 - \frac{1}{6} M_{ii} + \frac{1}{12} F_{ij}^2 \right).\end{aligned}\quad (10)$$

In these formulas $E_i = F_{0i}$ is the electric field and we use the convenient notation $M_{\mu\nu} = \mathcal{D}_\mu \mathcal{D}_\nu M$, $E_{\mu i} = \mathcal{D}_\mu E_i$, etc. The quantities φ_n are functions of $\Omega(x)$ and the ratio τ/β^2 , namely

$$\varphi_n(\Omega) = \frac{(4\pi\tau)^{1/2}}{\beta} \sum_{p_0} \tau^{n/2} Q^n e^{\tau Q^2}. \quad (11)$$

Here, Q given in (9) and p_0 runs over the Matsubara frequencies, $2\pi k/\beta$ (k integer) for bosons and $2\pi(k + \frac{1}{2})/\beta$ for fermions. Consequently, there are bosonic and fermionic versions for each function φ_n , related through

the replacement $\Omega \rightarrow -\Omega$. The φ_n 's are related to the special functions θ_3 and θ_4 and their derivatives [31].

The first thing to note is that, due to the summation over p_0 , the $\varphi_n(\Omega)$'s are one-valued functions of Ω and so gauge invariance is preserved. (The quantity $\log \Omega$ is covariant under small gauge transformations but many-valued under large ones.) Actually, if we use Poisson's summation formula for, say, the function φ_0 we obtain

$$\varphi_0(\Omega) = \sum_{k \in \mathbb{Z}} \Omega^k e^{-k^2 \beta^2 / 4\tau} \quad (\text{bosonic}). \quad (12)$$

Second, we note that the expansion contains coefficients of half-integer order with local operators of odd dimension. Due to the property $\varphi_n(\Omega) = (-1)^n \varphi_n(\Omega^{-1})$ the symmetry under $D_0 \rightarrow -D_0$, which is manifest in the Klein-Gordon operator, is preserved. Odd dimensional terms vanish at zero temperature except in presence of boundaries [32]. One of the techniques to deal with boundaries is the method of images. The path integral method suggests that the presence of these terms at finite temperature can also be understood from the fact that the particle sees its own image from its periodic replicas.

In the zero temperature limit the functions φ_n become constant. For even n , they take the values $(-\frac{1}{2})^{n/2} (n-1)!!$ and vanish for odd n . It is easy to verify that the thermal coefficients reproduce the zero temperature ones in this limit and Lorentz invariance is restored. As illustrated in (12) the low temperature or low τ corrections are exponentially suppressed in a_n^T .

The thermal coefficients have been collected according to the mass dimension of the local operators. At zero temperature this is equivalent to an expansion in powers of τ , but at finite temperature it corresponds to expand in powers of τ keeping τ/β^2 and Ω constant. Due to the exponential suppression, a strict expansion in τ for fixed β would yield just the zero temperature result. (This accounts for the well known fact that ultraviolet divergences and anomalies are temperature independent [33].)

Also useful is the expansion of the trace of the heat kernel operator $\text{Tr}(e^{-\tau K}) = \sum_n \tau^n \int d^d x \text{tr}(b_n^T)$. To mass dimension six we find

$$\begin{aligned}b_0^T &= \varphi_0 \\ b_1^T &= -\varphi_0 M \\ b_2^T &= \frac{1}{2} \varphi_0 M^2 - \frac{1}{3} \varphi_2 E_i^2 + \frac{1}{12} \varphi_0 F_{ij}^2 \\ b_{5/2}^T &= -\frac{1}{6} \varphi_1 \{M_i, E_i\} \\ b_3^T &= -\frac{1}{6} \varphi_0 M^3 + \frac{1}{6} \varphi_2 M_0^2 - \frac{1}{12} \varphi_0 M_i^2 \\ &\quad + \frac{1}{3} \varphi_2 E_i M E_i - \left(\frac{1}{12} \varphi_0 + \frac{1}{3} \varphi_2 + \frac{2}{15} \varphi_4 \right) E_{0i}^2 \\ &\quad + \frac{1}{30} \varphi_2 E_{ii}^2 + \frac{1}{30} \varphi_2 F_{0ij}^2 - \frac{1}{15} \varphi_2 E_i F_{ij} E_j \\ &\quad - \frac{1}{12} \varphi_0 F_{ij} M F_{ij} - \frac{1}{60} \varphi_0 F_{ii}^2 + \frac{1}{90} \varphi_0 F_{ij} F_{jk} F_{ki}.\end{aligned}\quad (13)$$

As is well known, the expression for b_n^T is intrinsically not unique due to integration by parts and the trace cyclic property. At zero temperature these operations do not mix different orders, but at finite temperature this is no longer true since the commutators (in particular covariant derivatives) of $\Omega(x)$ generate dimensional terms [34]. This explains why, e.g. $b_{3/2}^T$ vanishes although $\text{Tr}(a_{3/2}^T)$ is not identically zero. Of course, the trace of the heat kernel itself is unique and the ambiguity amounts to a reorganization of the asymptotic expansion (no such ambiguity exists for the a_n^T). The heat kernel is symmetric under transposition $ABC \dots \rightarrow \dots CBA$ and this symmetry is manifest in the b_n^T (using the cyclic property). It can be noted that, exactly as in the zero temperature case [8], the coefficients a_n^T can be obtained from the b_n^T through functional derivation with respect to $M(x)$.

In addition to the explicit formulas, one of the interesting results of this calculation is of qualitative type, namely, the operators appearing in the coefficients a_n^T or b_n^T can be grouped according to the weights φ_k . This indicates a definite pattern of Lorentz symmetry breaking at finite temperature (even in the absence of scalar potential). This pattern will manifest itself in applications such as those to renormalization group flow at finite temperature [35], the general form of the chiral perturbation theory effective Lagrangian at finite temperature, etc.

The fundamental role played by the Polyakov loop is further illustrated by considering the trivially solvable case of a homogeneous relativistic quantum gas. In the absence of external fields (other than a mass m), the lowest order term $b_0^T = \varphi_0$ is sufficient to obtain the effective action and so the grand canonical potential ($e^{-\tau m^2}$ exactly factorizes out of the heat kernel) [38]. A simple integral transform to pass from the exponential to the logarithm function reproduces the standard result, with distribution function $(e^{\beta(\omega_k - \mu)} - 1)^{-1} - (e^{\beta(\omega_k + \mu)} - 1)^{-1}$ (for bosons) [36]. We want to emphasize the relation between the chemical potential μ and the Polyakov loop Ω . The chemical potential couples as a constant additive term in the scalar potential. Because it is constant, μ does not contribute to the local operators, since there $A_0(x)$ only appears through the covariant derivative $\mathcal{D}_0$. If the Polyakov loop were absent in the formulas, μ would not appear at all in the partition function. An obviously incorrect result. It is also noteworthy that the periodic dependence of the heat kernel on $\log \Omega$ implies the well known fact that the partition function is periodic in $\beta\mu$ with period $2\pi i$ (a consistency condition due to its coupling to the quantized charge operator). The Polyakov loop thus appears as a generalization of the factor $e^{\beta\mu}$ to non constant and non Abelian gauge fields. In curved space-times, in addition to the Polyakov loop of the gauge connection A_μ , there is a Polyakov loop tied to the parallel transport connection Γ_μ , with consequences in field theory in presence of gravitational fields.

This work is supported in part by funds provided by

the Spanish DGI with grant no. BFM2002-03218, and Junta de Andalucía grant no. FQM-225.

-
- [1] J.S. Schwinger, Phys. Rev. 82 (1951) 664,
 - [2] B.S. Dewitt, Phys. Rept. 19 (1975) 295,
 - [3] R.T. Seeley, Proc. Symp. Pure. Math. 10 (1967) 288,
 - [4] P.B. Gilkey, J. Diff. Geom. 10 (1975) 601,
 - [5] M. Atiyah, R. Bott and V.K. Patodi, Invent. Math. 19 (1973) 279,
 - [6] S.W. Hawking, Commun. Math. Phys. 55 (1977) 133,
 - [7] K. Fujikawa, Phys. Rev. D21 (1980) 2848,
 - [8] R.D. Ball, Phys. Rept. 182 (1989) 1,
 - [9] M. Bordag, U. Mohideen and V.M. Mostepanenko, Phys. Rept. 353 (2001) 1, quant-ph/0106045,
 - [10] A.A. Bytsenko et al., Phys. Rept. 266 (1996) 1,
 - [11] R. Camporesi, Phys. Rept. 196 (1990) 1,
 - [12] A.A. Bel'kov, A.V. Lanyov and A. Schaale, Comput. Phys. Commun. 95 (1996) 123, hep-ph/9506237,
 - [13] A.E.M. van de Ven, Class. Quant. Grav. 15 (1998) 2311, hep-th/9708152,
 - [14] I.G. Moss and W. Naylor, Class. Quant. Grav. 16 (1999) 2611, gr-qc/0101125,
 - [15] D. Fliegner et al., Annals Phys. 264 (1998) 51, hep-th/9707189,
 - [16] I.G. Avramidi, Nucl. Phys. B355 (1991) 712,
 - [17] V.P. Gusynin, Phys. Lett. B225 (1989) 233,
 - [18] N.P. Landsman and C.G. van Weert, Phys. Rept. 145 (1987) 141,
 - [19] A.A. Actor, Phys. Rev. D50 (1994) 6560,
 - [20] C. García-Recio and L.L. Salcedo, Phys. Rev. D63 (2001) 045016, arXiv:hep-th/0007183,
 - [21] S. Deser, L. Griguolo and D. Seminara, (2002), hep-th/0212140,
 - [22] T.T. Wu and C.N. Yang, Phys. Rev. D12 (1975) 3843,
 - [23] H. Boschi-Filho, C.P. Natividade and C. Farina, Phys. Rev. D45 (1992) 586,
 - [24] A.A. Actor and K.D. Rothe, J. Phys. A33 (2000) 4585, hep-th/0002208,
 - [25] D.Y. Xu and X.F. Xu, J. Phys. A26 (1993) 7521,
 - [26] N. Nakazawa and T. Fukuyama, Nucl. Phys. B252 (1985) 621,
 - [27] G. Dunne, K.Y. Lee and C.h. Lu, Phys. Rev. Lett. 78 (1997) 3434, hep-th/9612194,
 - [28] L.L. Salcedo, Phys. Lett. B530 (2002) 244, hep-th/0201057,
 - [29] L.L. Salcedo and E. Ruiz Arriola, Ann. Phys. 250 (1996) 1, arXiv:hep-th/9412140,
 - [30] L.L. Salcedo, Nucl. Phys. B549 (1999) 98, arXiv:hep-th/9802071,
 - [31] M. Abramowitz and I.A. Stegun, Handbook of mathematical functions (Dover, New York, 1970),
 - [32] D.M. McAvity and H. Osborn, Class. Quant. Grav. 8 (1991) 603,
 - [33] L. Dolan and R. Jackiw, Phys. Rev. D9 (1974) 3320,
 - [34] E. Megías, E. Ruiz Arriola and L.L. Salcedo, (2003), in preparation.
 - [35] B.J. Schaefer and H.J. Pirner, Nucl. Phys. A660 (1999) 439, nucl-th/9903003,
 - [36] H.E. Haber and H.A. Weldon, Phys. Rev. Lett. 46 (1981) 1497,

[37] Further details, as well as the rather long coefficients $a_{5/2}^T$ and a_3^T , will be presented elsewhere [34].

[38] The effects of external fields can be included through higher orders in the heat kernel expansion.